

INTERACTIONS OF NEGATIVE K MESONS
IN FLIGHT IN NUCLEAR EMULSION

Frank Hunter Featherston

31

INTERACTIONS OF NEGATIVE K MESONS IN FLIGHT
IN NUCLEAR EMULSION

* * * * *

Frank Hunter Featherston

INTERACTIONS OF NEGATIVE K MESONS IN FLIGHT
IN NUCLEAR EMULSION

by
Frank Hunter Featherston

Submitted in partial fulfillment
of the requirements for the degree of

MASTER OF SCIENCE
IN
PHYSICS

United States Naval Postgraduate School
Monterey, California

1957

INTERACTIONS OF NEGATIVE K MESONS IN FLIGHT
IN NUCLEAR EMULSION

by

Frank Hunter Featherston

This work is accepted as fulfilling
the thesis requirements for the degree of

MASTER OF SCIENCE

IN

PHYSICS

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

INTERACTIONS OF NEGATIVE K MESONS IN FLIGHT
IN NUCLEAR EMULSION

Frank Hunter Featherston

(M. S. Thesis)

April 30, 1957

Submitted in partial fulfillment
of the requirements for the degree of

MASTER OF SCIENCE

in

PHYSICS

United States Naval Postgraduate School
Monterey, California

Printed for the U. S. Atomic Energy Commission

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contact No. W-7193-eng-48

INTERACTIONS OF NEGATIVE IONS WITH
POLYMER EMULSION

Paul H. Plesch

(M. S. Thesis)

April 10, 1952

Submitted in partial fulfillment
of the requirements for the degree of

MASTER OF SCIENCE

in

PHYSICS

United States Naval Postgraduate School
Monterey, California

Approved for the U. S. Atomic Energy Commission

INTERACTIONS OF NEGATIVE K MESONS IN FLIGHT IN NUCLEAR EMULSION

Frank Hunter Featherston

Radiation Laboratory
University of California
Berkeley, California

April 30, 1957

ABSTRACT

A survey of interactions of K^- mesons in flight in nuclear emulsion has been made. Seventy-seven in-flight interactions, seven decays in flight, and four inelastic scatterings were found.

A K^- meson mean free path in nuclear emulsion of 30.0 ± 3.0 cm has been calculated.

A K^- -meson lifetime of $1.31_{-.28}^{+.38} \times 10^{-8}$ sec has been calculated.

All interactions were consistent with the conservation of strangeness.

A general qualitative comparison is made between the in-flight and at-rest interactions of K^- mesons.

Two decays in flight were measured, one leading to a definite establishment of a $K^-_{\pi 2}$ decay mode, the other to a strong suggestion of a $K^-_{\mu 2}$ mode.

The angular distribution of the decay pions from 37 Σ hyperons produced by K^- interactions is reported. The distribution tends towards isotropy.

SYNTHESIS OF POLYMERIZABLE MONOMERS IN FLUORINE

Yoshio Imai, Tetsuo Imai

Department of Chemistry
University of California
Berkeley, California

April 19, 1957

ABSTRACT

A series of monomers of K. monomers in light in solution
monomers in solution. The monomers in light in solution
monomers in light, and the monomers in light in solution
K. monomers in light in solution of 20.5-1.0 g
has been calculated.

5.5 monomers in light in solution of 20.5-1.0 g
monomers in light in solution of 20.5-1.0 g

monomers

A series of monomers in light in solution of 20.5-1.0 g
monomers in light in solution of 20.5-1.0 g

Two monomers in light in solution of 20.5-1.0 g
monomers in light in solution of 20.5-1.0 g

The monomers in light in solution of 20.5-1.0 g
monomers in light in solution of 20.5-1.0 g

TABLE OF CONTENTS

Section	Title	Page
I	Introduction	
	The K Meson	1
	Strangeness	3
	Nuclear Emulsions as a Research Tool	6
II	General Experimental Method	
	Exposure and Development	9
	Scanning and Measurement	10
III	Results	
	Summary	12
	Mean Free Path of K^- Mesons in Emulsion	14
	Mean Lifetime of the K^- Meson	14
	Interaction Considerations	15
	Comparison with K^+ Mesons	16
	Interactions in Flight versus Absorptions at Rest	17
	Prong Distribution	17
	Effects of K^- -Meson Kinetic Energy	19
	Total Visible Energy Spectrum	21
	The Pi-Meson Spectrum	23
	Decays in Flight.	26
	K^- Meson Inelastic Scattering	28
	$K^- + H$ Compilation	29
	K^- Interactions in Flight: Conclusions	29
	Σ Decay Spectrum	29
	Σ^+ Branching Ratio	30
	Acknowledgments	33
	References	34

LIST OF ILLUSTRATIONS AND TABLES

Figure		Page
1.	Prong distributions for K^- -meson events at rest and in flight	18
2.	Comparison of sums of kinetic energies of associated Σ and π	20
3.	Comparison of total visible energy release.	22
4.	Pi-meson spectrum from K^- absorption stars	24
5.	Kinetic-energy spectrum of π mesons from K^- interactions in flight	25
6.	K^- -meson decays: momentum of the secondary particle as a function of K^- -meson momentum (for a laboratory angle of 58.0°)	27
7.	Angular spectrum of decay pions from hyperons produced in K^- -absorption stars	31
Tables		
I.	Length of K^- Meson Track Scanned in Energy Intervals.	13
II.	Results from inelastic scattering of K^- mesons	28

LIST OF CONTENTS

Page	Topic	Page	Topic
1	1. The nature of the problem	1	1. The nature of the problem
2	2. The scope of the problem	2	2. The scope of the problem
3	3. The method of solution	3	3. The method of solution
4	4. The results of the solution	4	4. The results of the solution
5	5. The conclusions of the solution	5	5. The conclusions of the solution
6	6. The bibliography	6	6. The bibliography
7	7. The index	7	7. The index
8	8. The appendix	8	8. The appendix
9	9. The list of figures	9	9. The list of figures
10	10. The list of tables	10	10. The list of tables
11	11. The list of references	11	11. The list of references
12	12. The list of symbols	12	12. The list of symbols
13	13. The list of abbreviations	13	13. The list of abbreviations
14	14. The list of units	14	14. The list of units
15	15. The list of definitions	15	15. The list of definitions
16	16. The list of formulas	16	16. The list of formulas
17	17. The list of equations	17	17. The list of equations
18	18. The list of theorems	18	18. The list of theorems
19	19. The list of lemmas	19	19. The list of lemmas
20	20. The list of propositions	20	20. The list of propositions
21	21. The list of corollaries	21	21. The list of corollaries
22	22. The list of examples	22	22. The list of examples
23	23. The list of exercises	23	23. The list of exercises
24	24. The list of problems	24	24. The list of problems
25	25. The list of projects	25	25. The list of projects
26	26. The list of assignments	26	26. The list of assignments
27	27. The list of tests	27	27. The list of tests
28	28. The list of quizzes	28	28. The list of quizzes
29	29. The list of exams	29	29. The list of exams
30	30. The list of papers	30	30. The list of papers
31	31. The list of theses	31	31. The list of theses
32	32. The list of dissertations	32	32. The list of dissertations
33	33. The list of books	33	33. The list of books
34	34. The list of journals	34	34. The list of journals
35	35. The list of magazines	35	35. The list of magazines
36	36. The list of newspapers	36	36. The list of newspapers
37	37. The list of radio programs	37	37. The list of radio programs
38	38. The list of television programs	38	38. The list of television programs
39	39. The list of films	39	39. The list of films
40	40. The list of records	40	40. The list of records
41	41. The list of tapes	41	41. The list of tapes
42	42. The list of CDs	42	42. The list of CDs
43	43. The list of DVDs	43	43. The list of DVDs
44	44. The list of Blu-rays	44	44. The list of Blu-rays
45	45. The list of video games	45	45. The list of video games
46	46. The list of computer games	46	46. The list of computer games
47	47. The list of board games	47	47. The list of board games
48	48. The list of card games	48	48. The list of card games
49	49. The list of dice games	49	49. The list of dice games
50	50. The list of puzzles	50	50. The list of puzzles
51	51. The list of riddles	51	51. The list of riddles
52	52. The list of jokes	52	52. The list of jokes
53	53. The list of cartoons	53	53. The list of cartoons
54	54. The list of comics	54	54. The list of comics
55	55. The list of novels	55	55. The list of novels
56	56. The list of short stories	56	56. The list of short stories
57	57. The list of plays	57	57. The list of plays
58	58. The list of operas	58	58. The list of operas
59	59. The list of musicals	59	59. The list of musicals
60	60. The list of ballets	60	60. The list of ballets
61	61. The list of operettas	61	61. The list of operettas
62	62. The list of pantomimes	62	62. The list of pantomimes
63	63. The list of comedies	63	63. The list of comedies
64	64. The list of tragedies	64	64. The list of tragedies
65	65. The list of dramas	65	65. The list of dramas
66	66. The list of farces	66	66. The list of farces
67	67. The list of satires	67	67. The list of satires
68	68. The list of romances	68	68. The list of romances
69	69. The list of epics	69	69. The list of epics
70	70. The list of legends	70	70. The list of legends
71	71. The list of myths	71	71. The list of myths
72	72. The list of fairy tales	72	72. The list of fairy tales
73	73. The list of fables	73	73. The list of fables
74	74. The list of parables	74	74. The list of parables
75	75. The list of allegories	75	75. The list of allegories
76	76. The list of metaphors	76	76. The list of metaphors
77	77. The list of similes	77	77. The list of similes
78	78. The list of personifications	78	78. The list of personifications
79	79. The list of hyperboles	79	79. The list of hyperboles
80	80. The list of oxymorons	80	80. The list of oxymorons
81	81. The list of paradoxes	81	81. The list of paradoxes
82	82. The list of ironies	82	82. The list of ironies
83	83. The list of sarcasms	83	83. The list of sarcasms
84	84. The list of puns	84	84. The list of puns
85	85. The list of wordplay	85	85. The list of wordplay
86	86. The list of anagrams	86	86. The list of anagrams
87	87. The list of palindromes	87	87. The list of palindromes
88	88. The list of acrostics	88	88. The list of acrostics
89	89. The list of alliterations	89	89. The list of alliterations
90	90. The list of onomatopoeias	90	90. The list of onomatopoeias
91	91. The list of personifications	91	91. The list of personifications
92	92. The list of metaphors	92	92. The list of metaphors
93	93. The list of similes	93	93. The list of similes
94	94. The list of personifications	94	94. The list of personifications
95	95. The list of hyperboles	95	95. The list of hyperboles
96	96. The list of oxymorons	96	96. The list of oxymorons
97	97. The list of paradoxes	97	97. The list of paradoxes
98	98. The list of ironies	98	98. The list of ironies
99	99. The list of sarcasms	99	99. The list of sarcasms
100	100. The list of puns	100	100. The list of puns

Out of a welter of early results and symbols, the heavy mesons have come to be known as the K mesons. This assignment of a definite name to these mesons by no means connotes the same unanimity and simplicity of characteristics and properties as are associated with the older well-established particles. In truth, within the realm of K mesons, it has been found necessary to present a more precise listing of these particles, based first, of course, upon electric charge, but more importantly, upon the several ways in which the K meson can decay (more properly, a phenomenological classification).

On this basis the so-called heavy or K mesons, can be listed as follows, with the decay modes indicated [1, 2]:

$$K^+_{\pi 3}, \tau^+$$

$$K^+_{\mu 2}$$

$$K^+_{\pi 2}, \theta^+$$

$$K^+_{\mu 3}$$

$$K^+_{e 3}$$

$$K^+ \rightarrow \pi^+ + \pi^- + \pi^+$$

$$K^+ \rightarrow \pi^+ + \pi^0 + \pi^0$$

$$K^+ \rightarrow \mu^+ + \nu$$

$$K^+ \rightarrow \pi^+ + \pi^0$$

$$K^+ \rightarrow \mu^+ + \pi^0 + \nu$$

$$K^+ \rightarrow e^+ + ? + ?$$

$$\theta^0, \theta^0_1 (?)$$

$$\theta^0, \theta^0_2 (?)$$

$$K^0 \rightarrow \pi^+ + \pi^-$$

$$(K^0 \rightarrow \pi^0 + \pi^0)$$

$$K^0 \rightarrow \pi^\pm + e^\mp + \nu$$

$$K^0 \rightarrow \pi^\pm + \mu^\mp + \nu (?)$$

$$K^-_{\pi 2}$$

$$K^-_{\mu 2}$$

$$K^-_{\mu 3}$$

$$K^-_{e 3}$$

$$K^-_{\pi 3}, \tau^-$$

$$K^- \rightarrow \pi^- + \pi^0$$

$$(K^- \rightarrow \mu^- + \nu)$$

$$(K^- \rightarrow \mu^- + \pi^0 + \nu)$$

$$K^- \rightarrow e^- + (?) + (?)$$

$$K^- \rightarrow \pi^- + \pi^- + \pi^+$$

$$(K^- \rightarrow \pi^- + \pi^0 + \pi^0)$$

Out of a welter of early results and symbols, the heavy mesons have come to be known as the K mesons. This assignment of a definite name to these mesons by no means connotes the same unanimity and simplicity of characteristics and properties as are associated with the older well-established particles. In truth, within the realm of K mesons, it has been found necessary to present a more precise listing of these particles, based first, of course, upon electric charge, but more importantly, upon the several ways in which the K meson can decay (more properly, a phenomenological classification).

On this basis the so-called heavy or K mesons, can be listed as follows, with the decay modes indicated [1, 2]:

$K^+_{\pi 3}, \tau^+$	$K^+ \rightarrow \pi^+ + \pi^- + \pi^+$
$\tau^+_{\pi 3}, \tau^+$	$K^+ \rightarrow \pi^+ + \pi^0 + \pi^0$
$K^+_{\mu 2}$	$K^+ \rightarrow \mu^+ + \nu$
$K^+_{\pi 2}, \theta^+$	$K^+ \rightarrow \pi^+ + \pi^0$
$K^+_{\mu 3}$	$K^+ \rightarrow \mu^+ + \pi^0 + \nu$
$K^+_{e 3}$	$K^+ \rightarrow e^+ + ? + ?$
$\theta^0, \theta^0_1 (?)$	$K^0 \rightarrow \pi^+ + \pi^-$
$\theta^0, \theta^0_2 (?)$	$(K^0 \rightarrow \pi^0 + \pi^0)$
	$K^0 \rightarrow \pi^\pm + e^\mp + \nu$
	$K^0 \rightarrow \pi^\pm + \mu^\mp + \nu (?)$
$K^-_{\pi 2}$	$K^- \rightarrow \pi^- + \pi^0$
$K^-_{\mu 2}$	$(K^- \rightarrow \mu^- + \nu)$
$K^-_{\mu 3}$	$(K^- \rightarrow \mu^- + \pi^0 + \nu)$
$K^-_{e 3}$	$K^- \rightarrow e^- + (?) + (?)$
$K^-_{\pi 3}, \tau^-$	$K^- \rightarrow \pi^- + \pi^- + \pi^+$
	$(K^- \rightarrow \pi^- + \pi^0 + \pi^0)$

In this listing, the symbol shown on the left of each decay mode is the presently accepted code by which each of the decay modes is designated. The " τ " and " θ " symbols have persisted from the earlier and original designations of these particular decay modes of the K meson. Those decay modes in parentheses have not actually been observed, but are to be expected from charge symmetry.

Strangeness

From this formidable listing of K-meson decay modes, one could quite properly ask whether we are witnessing the decay of two or more different particle species or whether we are seeing the several decay modes of a single species of particle, as the "K" designation implies.

This first question of the multiplicity of decay of the K meson is quickly joined by another experimental puzzle. Both hyperons and heavy mesons are abundantly produced in high-energy nuclear interactions, which means copious production on a time scale of the order of 10^{-23} sec; yet the lives of these particles, once they are produced, are very long by comparison ($\sim 10^{-8}$ to $\sim 10^{-10}$ sec). This discrepancy of approximately 10^{13} units of time scale is a seeming contradiction of reversibility principles, by which one would expect the decay of these particles in something akin to 10^{-23} second.

A suitable explanation of these effects began to take form in 1952 when Pais proposed that heavy mesons and hyperons can be produced only in pairs, i. e., in association with each other [3]. This associated production scheme indicated that there are certain selection rules that prevent strong interactions from being operative as part of the decay mechanism.

Finally in 1954, Gell-Mann, Nishijima, and others put forth the hypothesis that explained the apparent dilemma [4]. They

introduced a new property of matter, "strangeness," and stipulated certain selection rules based upon the assignment of a new quantum number to each of the particles in a process (this new number itself they called "strangeness", S). Certain properties of this new number were that it (1) must be integral, (2) could be either positive or negative, and (3) would be a constant of motion in strong interactions, but would change by ± 1 in weak interactions.

From this scheme of things, particle-interaction processes were divided into three classes:

a. Fast Interactions (i.e., direct production)

- (1) time scale $\sim 10^{-23}$ sec.
- (2) satisfaction of all conservation laws (energy, momentum, angular momentum, parity, charge, heavy particle number, and strangeness). Thus, $\Delta S = 0$.
- (3) process does not include emission of a gamma ray.

b. Electromagnetic Interactions

- (1) involve the emission of a gamma ray.
- (2) weaker by $1/137$ than the fast reactions
- (3) proceed on a time scale of $\sim 10^{-17}$ sec.

c. Slow Interactions

- (1) decays
- (2) proceed on a time scale of 10^{-10} sec.
- (3) can violate parity considerations
- (4) $\Delta S = \pm 1$

Thus, (a) would be the strong interactions which involve baryons, antibaryons, and mesons. We are here in the region of fully nuclear forces, in which mesons and hyperons are produced in high-energy nuclear interactions. The electromagnetic interaction (b) is the natural mechanism whereby the photon involves itself with all charged particles.

The slow (or weak) interactions (c) are the beta-decay process, the various heavy-meson and hyperon decays, and μ -mesonic absorption and decay.

The assignment of strangeness quantum numbers to the various particles is as follows:

$S = 0:$	$p, n; \pi^+, \pi^0, \pi^-$
$S = 1:$	K^0, K^+
$S = -1:$	$K^-, \bar{K}^0; \Lambda^0; \Sigma^+, \Sigma^0, \Sigma^-$
$S = -2:$	Ξ^0, Ξ^-

It is well to remember that two charge-conjugate particles must have equal and opposite values of S , because in a fast interaction a particle can be transposed to the other side of the reaction and become its own antiparticle without changing the fast nature of the reaction.

One finds the most direct experimental support of strangeness from a consideration of the charge degeneracy of strange-particle states and the z th component of the isotopic spin. From π -meson physics there comes an expression relating these:

$$Q - I_z - N/2 = 0,$$

where Q is the charge,

I_z is the third component of the isotopic spin,

N is the number of nucleons.

If, through some mechanism in strange-particle interactions, the conservation of isotopic spin remains but the above relation does not hold, i. e.,

$$Q - I_z - N/2 \neq 0,$$

then this sum must still be a quantity that is additively conserved, since Q , I_z , and N are all conserved separately. Thus, the basis of the strangeness scheme is a postulation that a new quantum number,

$$S = 2(Q - I_z - N/2),$$

can correlate and explain the strange-particle interactions.

An illustrative test of this is readily found in the Σ hyperons. The Σ^- has $S = -1$, $Q = -1$, and $N = 1$. I_z thus equals -1 . This indicates that there must be two additional particles of mass similar to that of the Σ^- , with $I_z = 0$ and 1 , and $Q = 0$ and $+1$, respectively. Experiment has found two such particles satisfying these conditions, Σ^0 and Σ^+ .

The consistency of its explanations has been the remarkable feature of the strangeness concept. There are no known violations of the strangeness-selection rules. Associated production, too, appears to be the order of business. As an explanation, the scheme has done yeomen's service in pointing the way through the dense underbrush of much contemporary research.

Nuclear Emulsions as a Research Tool

The techniques of high-energy particle research are necessarily of a very specialized nature. There are four general experimental systems used in high-energy particle research,

- (1) nuclear photographic emulsions,
- (2) cloud chambers,
- (3) bubble chambers,
- (4) scintillating and electronic counting networks.

Each of the four systems has both advantages and limitations peculiar to itself. Employment of a given system in a specific application may enhance or diminish certain of its capabilities.

As a stopping medium, nuclear emulsions are the heaviest by far of the investigative media, having a density of about 3.85 g/cc (average density of Ilford G. 5 emulsion) [5]. The basic elemental constituents of such an emulsion, with the corresponding approximate percentage compositions, by weight, are [5]:

Silver	47.5%	Hydrogen	1.4%
Bromine	35.0	Oxygen	6.5
Iodine	0.3	Nitrogen	1.9
Carbon	7.2	Sulfur	0.2

As a research tool, the emulsion provides a means for direct visual survey and measurement of the tracks left by ionizing particles as they pass through the emulsion stack. (Neutral particles of course, leave no tracks.)

A particle entering an emulsion may

- (1) Suffer energy degradation and slowing down, but succeed in leaving the emulsion; or
- (2) interact (or annihilate) with one of the constituent elemental nuclei while in flight; or
- (3) while in flight in the emulsion, decay, with the resulting particle(s) continuing on in the emulsion; or
- (4) be progressively slowed down until it comes to rest in the emulsion. Characteristically, then, the particle can either (if negative) be absorbed into a constituent nucleus, or (if positive) decay.

With a proper selection of physical parameters, an emulsion stack provides a wonderfully compact and portable means for observing the in-flight and terminal behavior of ionizing particles. Also, because of its great stopping power, the nuclear emulsion is a useful device for coping with short-lived particles when one is attempting to obtain a

measure of mean lifetime of the order of 10^{-8} to 10^{-13} sec. This is made possible through the greater density of the medium, which causes many of the particles to come to rest before they decay.

As one of the oldest of high-energy particles research systems, the art of employing nuclear emulsions has given rise to many refined auxiliary techniques. Through the use of these techniques one is able to

- (1) make an exhaustive analysis of specific and unusual events,
- (2) make very accurate mass determinations,
- (3) accomplish ready and reliable identification of particles,
- (4) precisely predict the response of the system to ionizing particles of any mass (this makes the emulsion well suited for exploratory detection attempts [6]),
- (5) make accurate ionization measurements.

The one important factor to appreciate when considering nuclear emulsion results is that the interacting nuclei of the emulsion proper are of several types, covering a wide range of atomic numbers. Silver and bromine nuclei are by far the most numerous and the heaviest, with an average A of 94. Hence, any discussion of emulsion results will be conditioned by this fact. This is not, as might be hastily inferred, a completely deleterious effect; for as pleasant-appearing as the simpler interaction phenomena of the bubble chambers may appear, particle interactions in complex nuclei have a definite appeal of their own.

The first of these is the fact that the
 second of these is the fact that the
 third of these is the fact that the
 fourth of these is the fact that the
 fifth of these is the fact that the
 sixth of these is the fact that the
 seventh of these is the fact that the
 eighth of these is the fact that the
 ninth of these is the fact that the
 tenth of these is the fact that the

- (1) The first of these is the fact that the
- (2) The second of these is the fact that the
- (3) The third of these is the fact that the
- (4) The fourth of these is the fact that the
- (5) The fifth of these is the fact that the
- (6) The sixth of these is the fact that the
- (7) The seventh of these is the fact that the
- (8) The eighth of these is the fact that the
- (9) The ninth of these is the fact that the
- (10) The tenth of these is the fact that the

The first of these is the fact that the
 second of these is the fact that the
 third of these is the fact that the
 fourth of these is the fact that the
 fifth of these is the fact that the
 sixth of these is the fact that the
 seventh of these is the fact that the
 eighth of these is the fact that the
 ninth of these is the fact that the
 tenth of these is the fact that the

The first of these is the fact that the
 second of these is the fact that the
 third of these is the fact that the
 fourth of these is the fact that the
 fifth of these is the fact that the
 sixth of these is the fact that the
 seventh of these is the fact that the
 eighth of these is the fact that the
 ninth of these is the fact that the
 tenth of these is the fact that the

II. GENERAL EXPERIMENTAL METHOD

Exposure and Development

This comprehensive study of a group of K^- mesons in flight interactions in nuclear emulsion is a modest (but typical) example of the directed application of research effort, made possible by the availability of relatively intense beams of artificially produced K mesons from the Bevatron.

In this instance an emulsion stack of Ilford G. 5 stripped nuclear emulsion, composed of 108 pellicles, each 4 by 7 inches and 600 microns thick, was exposed to a momentum-selected K^- beam of 420 Mev/c. Momentum selection was accomplished by utilizing the magnetic field of the Bevatron itself; quadrupole lenses were used to increase the intensity of the beam. This exposure was the cooperative effort of Gerson and Sulamith Goldhaber and Warren W. Chupp at the Berkeley Bevatron.

The beam to which the stack was exposed contained a mixture of both negative π mesons and the desired K^- mesons (in the approximate ratio of 3000 to 1). The π mesons presented ionization tracks of near-minimum intensity in the emulsion at this particular momentum; hence, they provided a convenient built-in comparative measure of ionization.

The K^- beam was emitted at 0° relative to the target. The beam was lead undeviated through the quadrupole focusing system, which meant that a background of contamination could appear in the exposure because of spurious products produced by portions of the beam striking the yokes of the focusing magnets. To remove this source of contamination, a sweeping magnet was placed behind the quadrupole system to remove all positive particles (i. e., protons).

The emulsion stack was aligned for exposure so that its long dimension was longitudinally placed in the beam. Thus, the particle tracks were parallel to the emulsion layers; the mean range of K^- mesons of this momentum (10.2 cm) was, conveniently, about two-thirds the length of the plate.

Development of the stack followed a modified "Bristol" procedure. Subsequent location of the "x" and "y" coordinates of a given position within each plate-mounted pellicle was facilitated by the contact printing of a grid system of coordinates on the surface of the emulsion, with numbered grids occurring every millimeter [7].

Scanning and Measurement

Inspection of the plate-mounted and developed layers of a photographic nuclear emulsion is accomplished with high-resolution microscopes. This "scanning" can be executed by following either of two general techniques: (1) area scanning, in which the observer looks at all events within a given area of the plate, looking for interaction stars, and subsequently following the causative particle back from the event to where it entered the emulsion; (2) along-the-track scanning, in which scanning is done transversely across the beam of entering particle tracks near the edge of the plate. On the readily determined basis of their relative degree of ionization, tracks of the appropriate ionization can be followed and their terminal behavior recorded.

Along-the-track scanning introduces a minimum of bias towards finding or rejecting any particular species of terminal behavior. All types of track endings are seen with equal facility, since all tracks are followed from where they enter the emulsion, before an event occurs. Area scanning, on the other hand, must--by the nature of the technique and from the inevitable human considerations involved--result in not seeing all the events with no secondaries or with light,

minimal secondaries. Area scanning has its place under certain circumstances (cosmic-ray exposures, or surveys for particular types of interactions or interaction products, for example).

In the exposure study made in gathering the data for this report, transverse scanning near the exposed edge of the plates readily revealed those particles with a degree of ionization approximately 1.8 times minimum, the theoretically predictable degree of ionization characteristic of K^- mesons of the selected momentum in Ilford G.5 emulsion.

Each track so selected was followed to its end in the emulsion. Every reported event was unambiguously identified as being caused by a K^- meson, either by successive grain counts (ionization measurements) if sufficient track length was available, or by multiple-scattering techniques, utilizing the constant-sagitta method, to obtain a measure of the particle's $p\beta$, to be compared with its relative ionization.

All interaction products were followed either to their endings or until they left the stack. All prongs not identifiable as electrons, mesons, hyperons, or hyperfragments were termed "stable charged" prongs and were assigned kinetic energies from their measured ranges upon the assumption of a protonic mass. This leads to a resulting underestimate of the kinetic energy of approximately one third of the particles, which are actually deuterons, tritons, alpha particles, or recoil fragments. All prongs less than 5 microns long were termed "recoils" and were not considered in either prong or energy distributions.


III. RESULTS

Summary

All together, seventy-seven interactions in flight of K^- mesons have been identified. Seven additional events were identified as decays in flight of K^- mesons; four other events have been established as representing the inelastic scattering of K^- mesons by the constituent nuclei of the emulsion.

Of the 77 in-flight events, 13 produced charged Σ hyperons, 10 of which had visible associated π mesons. Three events produced charged Σ hyperons without any visible π mesons. Similarly, 12 events produced π mesons without identifiable Σ hyperons. Thus, 17% of the event produced identifiable Σ hyperons; 29% of the events produced π mesons; 13% of the events produced visible Σ hyperons and π mesons in association.

Of the 13 Σ 's seen, two were Σ^- , five were Σ^+ , and six decayed in flight and hence could be either Σ^+ or Σ^- . Of these six, one could be considered Σ^- because the Σ itself was the sole visible reaction product; thus, there is a possibility that the interaction went via the available channel:



One star produced a hyperfragment, which decayed nonmesically.

One of the ten observed $\Sigma + \pi$ events was established as representing the capture of a K^- meson in hydrogen [8].

Forty-eight of the events produced only charged stable, i. e., "evaporation", prongs.

Three of the K^- tracks ended in flight with no visible prongs. These are classified as disappearances and are included in the total of 77.

These results represent a scanning effort of 25.71 meters of K^- meson track (in which, in addition, 226 K^- -meson absorptions at rest have been identified)[9]. This figure for track length includes two corrective factors: (1) The last 2 mm of each track that comes to rest (corresponding to the last 16 Mev of kinetic energy) have been omitted from consideration because a decay in flight in this region is difficult to identify; (2) The first 5 mm of each track have not been included because the primary particle causing an event in this early region cannot be easily identified as a K particle. Table I is a tabulation of the length of track scanned in the indicated energy intervals.

TABLE I

<u>Length of K^- Meson Track Scanned in Energy Intervals</u>	
<u>Energy Interval</u> (Mev)	<u>Track Length</u> (cm)
16 - 30	94.4
30 - 40	90.5
40 - 50	116.9
50 - 60	132.8
60 - 70	156.1
70 - 80	167.3
80 - 90	193.2
90 - 100	221.6
100 - 110	221.7
110 - 120	237.1
120 - 130	261.2
130 - 140	268.7
140 - 150	272.6
150 - 160	124.8
160 - 170	12.1
Total	25.71 meters

Mean Free Path of K^- Mesons in Nuclear Emulsion

When the track length and number of events of this report are combined with the corresponding data from Iloff [2], who reports 21 in-flight interactions in 4.90 meters of track, the mean free path in nuclear emulsion for K^- mesons is found to be

$$\lambda = 30.0 \pm 3.0 \text{ cm.}$$

This corresponds to a nuclear radius $R = (1.32 \pm 0.07 \times 10^{-13}) A^{1/3}$ cm.

Mean Lifetime of the K^- Meson

The mean lifetime of the K^- meson, τ_{K^-} , equals T/N , where T is the algebraic sum of the proper slowing-down times for each K^- track, calculated from where the particle is first registered to where it decays or interacts in flight, or up to a point 2 mm before it comes to rest. N is the total number of observed decays. T is calculated on the basis of the tables of Barkas and Young [10].

The results of this investigation show seven (7) decays in flight with a corresponding proper slowing-down time of 13.85×10^{-8} sec. Again, combining these results with those of Iloff et al. [11], who report 13 decays with a corresponding total proper slowing-down time of 12.4×10^{-8} sec, one obtains

$$\tau_{K^-} = 1.31^{+.38}_{-.28} \times 10^{-8} \text{ sec.}$$

(The error quoted is based on confidence limits for 68% probability for 20 events).

This mean life, of course, is a mixture of the several possible K^- decay modes. It is to be compared with

$$\tau_{K^-} = 1.49^{+.22}_{-.24} \times 10^{-8} \text{ sec,}$$

recently reported from the results of a counter experiment [12]. This counter-determined value is based on an experiment conducted four mean lives from the target; the value presented in this report is based on an investigation two mean lives from the target.

Interaction Considerations

The conservations of strangeness, charge, and heavy-particle number allow only the following one-nucleon interactions of K^- mesons at rest:

- (1) $K^- + p \rightarrow \Sigma^+ + \pi^-$,
- (2) $\quad \quad \rightarrow \Sigma^- + \pi^+$,
- (3) $\quad \quad \rightarrow \Sigma^0 + \pi^0$,
- (4) $\quad \quad \rightarrow \Lambda^0 + \pi^0$,
- (5) $K^- + n \rightarrow \Sigma^- + \pi^0$,
- (6) $\quad \quad \rightarrow \Sigma^0 + \pi^-$,
- (7) $\quad \quad \rightarrow \Lambda^0 + \pi^-$.

Certain types of two-pion reactions are allowed, but are negligible from phase-space considerations. Multiple nucleon capture also is allowed, but is considered to make only a small contribution ($\sim 10\%$) [9].

These interactions come about through the capture at rest of the K^- meson in a Bohr orbit of the atom, with subsequent absorption into the nucleus. These absorptions are strong ("fast") interactions, occurring in $\sim 10^{-23}$ sec. They must, of necessity, proceed with positive Q values.

For the $K^- + p$ reaction, (1) through (4), one must have clearly in mind the difference between capture of a K^- meson by a proton that is bound in the nucleus, and K^- capture by a free proton. The bound proton is moving in a nuclear potential distribution; consequently, when it captures a K^- meson, the reaction products must find their way free of the nucleus. Thus, their visible energies are modified by

their escapes. The parent nucleus is also left in an excited state and will emit so-called "evaporation" prongs.

By contrast, K^- capture by a free proton displays none of these complications. The full Q of the reaction is carried away in the primary reaction products. Kinematic resolution of these events is thus possible through accurate measurement of the secondary tracks.

Comparison with K^+ Mesons

In addition to the K^- -absorption interactions listed in (1) through (7), the K^- meson is permitted to interact with matter while in flight by two additional mechanisms.

- a. Scattering, as permitted by



This scattering corresponds to inelastic scattering from a nucleus, but in scattering from hydrogen, elastic scattering is also possible.

- b. Charge exchange, wherein



It is in these types of interaction, i. e., scattering and charge exchange, that positive K mesons display an exact correspondence to negative K -meson interaction types. Thus, the following K^+ meson interactions in flight are permitted:

- a. Scattering:



- b. Charge Exchange:



Because K^+ mesons have a strangeness value of +1, they are forbidden from undergoing absorption interactions corresponding to

their shapes. The present method is also not so tedious and will not require "specialized" groups.

By means of K , we obtain a new kind of distribution of the combinations. The total of the combinations is now in the form of a single system of points. Elements according to their values are possible through each of the members of the symmetry class.

Comparison with K elements

In addition to the K -combinations mentioned above in the text (1) the K members are found in constant with respect to (2) the by two additional members:

$$\begin{aligned} & \text{a. } \text{Symmetry, combinations:} \\ & \quad K^+ + K^- = K^0 \\ & \quad K^+ + K^- = K^0 \end{aligned}$$

Total symmetry combinations in the present system are: instead, but in addition to the combinations, which are already listed:

$$\begin{aligned} & \text{b. } \text{Change combinations:} \\ & \quad K^+ + K^- = K^0 \end{aligned}$$

It is in these cases of combinations: 1. a. symmetry and change combinations, that positive is added to the combinations and negative is added to the combinations. Thus, the combinations K^+ and K^- are added.

$$\begin{aligned} & \text{c. } \text{Change combinations:} \\ & \quad K^+ + K^- = K^0 \\ & \quad K^+ + K^- = K^0 \\ & \quad K^+ + K^- = K^0 \end{aligned}$$

By means of K members, the combinations are: 1. the same as the combinations mentioned above in the text, but in addition to the combinations, which are already listed:

the types (1) through (7) listed for K^- mesons. Thus, a K^+ meson that comes to rest in matter suffers Coulomb exclusion from the surrounding nuclei and can only decay (via any one of several decay modes).

Interactions in Flight versus Absorptions at Rest

In the published literature only a limited attempt has been made to compare K^- in flight behavior and K^- absorptions at rest [13]. This condition has been dictated by statistics, because in the energy region explored, the preponderance of K^- events in emulsion have been the absorptions of K^- mesons that have come to rest.

Although this report significantly increases the published statistics for in-flight interactions, the significance is still such as to permit only limited qualitative comparison. No doubt the immediate future will see the issuance of many data now in the incubation stage. Then, quantitative comparisons can be made.

Prong Distribution

Figure 1 is a comparison of the prong distributions produced by K^- absorptions at rest and K^- interactions in flight. The "at rest" portion is the published compilation of the 1956 Rochester Conference (449 events) [13]. The "in flight" portion represents an improvement of the Rochester data (where 56 events were reported from five separate research groups) by the addition of the 77 events of this report.

Two points can be made from this comparison:

- (1) The increase in \bar{N} , the average number of prongs per star, for stars in flight reflects an increase in nuclear excitation in these events.
- (2) There is justification for the premise that charge exchange is not a very large part of the K^- absorption cross section because there is no increase (in fact, there is a percentage decrease) in the number of stars with zero or one prong.

the same (1) through (11) under the "Measure" column. The same data
can be used for many other purposes. It can be used for
the purpose and can also be used for any other purpose.

Notes on the Table

In the present report, the data are arranged in two columns.
The first column is headed "Measure" and the second column is headed
"Value". The data are arranged in two columns. The first column is
headed "Measure" and the second column is headed "Value". The data
are arranged in two columns. The first column is headed "Measure" and
the second column is headed "Value".

Although the data are arranged in two columns, the data are
not arranged in two columns. The data are arranged in two columns.
The data are arranged in two columns. The data are arranged in two
columns. The data are arranged in two columns. The data are arranged
in two columns. The data are arranged in two columns. The data are
arranged in two columns. The data are arranged in two columns.

Table 1

Table 1 is a summary of the data. The data are arranged in
two columns. The first column is headed "Measure" and the second
column is headed "Value". The data are arranged in two columns.
The data are arranged in two columns. The data are arranged in two
columns. The data are arranged in two columns. The data are arranged
in two columns. The data are arranged in two columns. The data are
arranged in two columns. The data are arranged in two columns.

The data are arranged in two columns. The data are arranged in
two columns. The data are arranged in two columns. The data are
arranged in two columns. The data are arranged in two columns. The
data are arranged in two columns. The data are arranged in two
columns. The data are arranged in two columns. The data are arranged
in two columns. The data are arranged in two columns. The data are
arranged in two columns. The data are arranged in two columns.




Fig. 1. Prong distributions for K^- -meson events at rest and in flight.


Effects of K^- -Meson Kinetic Energy

The average kinetic energy at interaction of the 77 K^- mesons of this report was 105 Mev. Because one is looking for comparative effects between K^- interactions in flight and K^- absorptions at rest, an obvious effect is to see how (or whether) this 105 Mev asserts itself. The increase in \bar{N} in the previous prong distribution is doubtless one effect. Again, one must bear in mind the limited in-flight statistics.

Thirteen of the in-flight events produced single charged hyperons, and ten of these events exhibited an associated π meson. Only one of these ten constituted an interaction with a free proton; the other nine events occur with bound protons.

Figure 2 shows a comparison of the total kinetic energy of $\Sigma + \pi$ combinations for in-flight interactions and at-rest absorptions. The approximate mean energies of each distribution are, respectively, 140 and 80 Mev. The Q available from the general reaction $K^- + \text{nucleon} \rightarrow \Sigma + \pi$ is about 104 Mev. The "at rest" distribution forms a peak at about 80 Mev, which means that the Σ and π succeed in escaping the reaction volume with about 80 Mev of their original 104 Mev. The difference in energy goes into nuclear excitation, and possible inelastic scattering losses of the particles while still in the dense nuclear region.

The "in flight" portion of Fig. 2, then, would indicate that the primary reaction products carry off an additional 60 Mev of the now-available 105 Mev of kinetic energy, the remainder of this kinetic energy going into increased nuclear excitation (as indicated in the prong distribution, Fig. 1). The use of these specific energy figures is meant to convey the degree or apparent order of the effect seen, rather than to mean anything truly quantitative.



MU-13402

Fig. 2. Comparison of sums of kinetic energies of associated Σ and π .

Total Visible Energy Spectrum

Figure 3 is another comparative distribution, this time showing the total visible energy spectra. The following constitute the criteria for energy-value assignments:

(1) All charged stable prongs were assigned kinetic energies on the basis of an assumed protonic mass from their measured ranges. In each case, then, 8 Mev of binding energy was added (a minimum amount of energy necessary to remove a proton from the nucleus). As previously pointed out, this procedure actually sets a lower limit to approximately one third of this portion of the spectra.

(2) Pi mesons were assigned their measured kinetic energies plus a rest mass energy of 139.5 Mev.

(3) Hyperons were assigned their measured kinetic energies plus a Q value taken as the difference between the hyperon mass and the proton mass, i.e., 1189 Mev minus 938 Mev, which equals 251 Mev.

Bearing in mind, again, the difference in number of events at rest and in flight, one notices the following:

(a) The percentage decrease in the number of stars with less than 20 Mev. This is to be expected since it would take only a portion of the available 105 Mev of K^- meson kinetic energy going into increased nuclear excitation, to bring about this low energy shift.

(b) The identified $\Sigma + \pi$ events (cross-hatched) assert themselves at the extreme right of each distribution. The "in flight" events shift almost as a block, as previously indicated in Fig. 2.

One would expect some fraction, say about half, of the 105 Mev of average K^- kinetic energy to appear as a general translational effect along the entire length of the "in flight" distribution. Qualitatively, such an effect is observed.

Figure 1 shows the energy distribution of the system. The total energy is 1000 units.

The total energy is 1000 units. The energy is distributed among the system.

The energy is distributed among the system.

(1) The energy is distributed among the system.

(2) The energy is distributed among the system.

(3) The energy is distributed among the system.

(4) The energy is distributed among the system.

(5) The energy is distributed among the system.

(6) The energy is distributed among the system.

(7) The energy is distributed among the system.

(8) The energy is distributed among the system.

(9) The energy is distributed among the system.

(10) The energy is distributed among the system.

(11) The energy is distributed among the system.

(12) The energy is distributed among the system.

(13) The energy is distributed among the system.

(14) The energy is distributed among the system.

(15) The energy is distributed among the system.

(16) The energy is distributed among the system.

(17) The energy is distributed among the system.

(18) The energy is distributed among the system.

(19) The energy is distributed among the system.

(20) The energy is distributed among the system.


(21) The energy is distributed among the system.

(22) The energy is distributed among the system.

(23) The energy is distributed among the system.

(24) The energy is distributed among the system.

(25) The energy is distributed among the system.



MU-13403

Fig. 3. Comparison of total visible energy release.




Fig. 2. Comparison of two slightly different sections

The Pi-Meson Spectrum


Of considerable interest in the general study of K mesons is a determination of the frequency with which K^- mesons interact in the several available channels. A first approach would be to determine the $\Sigma^{\pm 0}/\Lambda^0$ production ratio. Because the Λ^0 hyperon will not register in the investigative medium, one must try to deduce this ratio from associated conditions that can be observed. This means that the observed charged-pion spectrum can be quite meaningful.

The statistics of the π -meson spectrum from K^- absorptions at rest are now quite good. Figure 4 shows a histogram of 169 such pions [9].

The π mesons produced in the free-nucleon capture of a K^- meson have characteristic kinetic energies of 90 or 150 Mev, depending upon whether the associated hyperon is a Σ or Λ . Figure 4 clearly displays a peak at 60 Mev both for the over-all distribution and for those pions produced in association with charged hyperons. This coincidence of peaks makes a very strong argument for the preferred production of energy-degraded 90-Mev π mesons produced in the $\Sigma + \pi$ reaction (as opposed to the $\Lambda + \pi$ type of reaction).

Figure 5 is a constant-area histogram showing the kinetic-energy distribution of the 22 charged pions observed from the 77 in-flight interactions of this report. The pion energies were determined by ranges where possible, or by grain count if the particle left the emulsion stack.

This spectrum of pions from the in-flight interactions is based on relatively poor statistics, but it does have one feature that subtly enhances the contention that Σ production is significantly preferred to Λ production; specifically, the pronounced high-energy peaking shows that higher-energy π mesons are not excluded from making their presence felt, if they are produced in the first place. This tends to



MU-13404

Fig. 4. Pi-meson spectrum from K^- absorption stars⁹.




Fig. 5. Kinetic-energy spectrum of π mesons from K^- interactions in flight. (Illustration is a constant-area histogram, reflecting the error in energy determination.)



Fig. 4. Histogram of energy levels of the α particles from the ^{210}Po source. The hatched bars represent the calculated energy levels and the solid bars represent the measured energy levels.

negate arguments that have said that there is a very strong bias against the emission of high-energy pions from complex-nuclei interactions (arguments based on the peculiarities of pion scattering inside the nucleus).


The inference, again, is that the 150-Mev pions of the $\Lambda + \pi$ reactions are infrequently produced. Thus the $\Lambda + \pi$ interaction channels for K^- mesons appear to be small.

Decays in Flight

In this study of K^- in-flight interactions, seven events were in-flight decays. An in-flight event was classified as a decay if it exhibited only one secondary prong, this prong having a degree of ionization less than that of the primary at the point of the decay. Such an event could represent an in-flight interaction with a single π -meson secondary. However, a survey of K^- absorptions at rest shows this to be a very infrequent star type (8 out of 325, in one survey) [2]. Since in-flight interactions have a much greater tendency to display evaporation prongs because of increased nuclear excitation, the frequency of occurrence of single π events is even further reduced.

Of the seven decays observed, two displayed secondaries that were amenable to measurement. One secondary was emitted at a space angle of 57.8° relative to the parent K^- meson. As Fig. 6 shows, this event (A) unambiguously resolves itself as representing the decay of a K^- meson via $K_{\pi 2}^- \rightarrow \pi^- + \pi^0$.

The second decay (B) occurred at a space angle of 57.5° relative to the K^- meson. Its kinematic resolution is not so well defined as is that of event A. Both events are displayed on Fig. 6, since they occur at nearly the same space angle. The angular error is very small ($\sim 1\%$), and this region of comparative momenta is fairly insensitive to small changes in space angle.



MU-13406

Fig. 6. K^- -meson decays: momentum of the secondary particle as a function of K^- -meson momentum (for a laboratory angle of 58.0°).




Fig. 1. Dependence of the relative residual amount of the polymer on the temperature of the polymerization. A - 100% (1000); B - 100% (1000).

Event B is more nearly commensurate with a $K^-_{\mu 2} \rightarrow \mu^- + \nu$ decay mode, as Fig. 6 indicates. When its resolution is attempted by use of a particle of π -mesonic mass, the fit to the necessary curve ($K^-_{\pi 2}$) is worse, further strengthening the $K^-_{\mu 2}$ conclusion.

K^- Meson Inelastic Scattering

Four inelastic scatterings of K^- mesons were observed. Table II summarizes the results.

TABLE II

Results from inelastic scattering of K^- mesons		
Kinetic energy of K^- meson at scatter (Mev)	Fractional energy loss, $\Delta E/E$	Space angle of scatter (degrees)
38	0.58	55.9
58	0.62	81.8
103	0.48	58.8
138	0.55	102.8

From isotopic spin considerations, the K^- ratio of (charge exchange/inelastic scattering) is ≤ 2 . Thus, the limit on the charge-exchange contribution to the absorption cross section is $\leq 8\%$ [9].

The large fractional energy loss ($\Delta E/E$) indicates that an attractive (negative) potential may be operating in K^- scattering. In addition to the four inelastic scatterings reported here, a recent summary of emulsion data [14] indicates seven other events, all with $\Delta E/E$ greater than 0.40. This effect is the opposite of that observed in the scattering of K^+ mesons. There the energy loss in inelastic collisions is small, $\Delta E/E \approx 0.35$, which can be ascribed to a repulsive potential [15].

K⁻ + H Compilation

For comparative purposes, a compilation of published results of K⁻ + H scattering and absorption is given below.

Group	Energy Interval (Mev)	Absorptions in flight	Number of scatters	Path length (meters)
Goldhaber et al. [9]	16 - 160	1	1	33
White et al. [16]	16 - 150	1	10	30
Barkas et al. [17]	30 - 90	$\frac{2}{4}$	$\frac{6}{17}$	$\frac{49.5}{112.5}$

From this the following K⁻ + H cross sections have been calculated:


- (1) Scattering 47^{+14}_{-11} mb.
- (2) Absorption 11^{+9}_{-5} mb.
- (3) Total 58^{+16}_{-13} mb.

K⁻ Interactions in Flight: Conclusions

Within the statistics of the data of this report, there is no apparent major difference between "in flight" interactions and "at rest" interactions of K⁻ mesons in nuclear emulsion. Interactions in flight display greater nuclear excitation, attributable to the extra available kinetic energy. The inelastic scattering data indicate the presence of an attractive nuclear potential.

Σ Decay Spectrum

The charged Σ hyperons can decay via



The angular distribution of these decay products with respect to the hyperon's in-flight direction can be used to investigate:

(1) Parity considerations. Lee and Yang [18] have shown that if the Σ has mixed parity, the decay spectrum should show a forward-backward asymmetry.

(2) Spin of the Σ hyperon. After the folding of such an angular distribution about 90° , one can get an indication of spin, independent of parity doublet considerations. Thus, an isotropic folded distribution would indicate a spin of $1/2$; any asymmetry would indicate a spin $> 1/2$ [19].

A survey has been made of the decays of Σ 's produced by K^- absorptions at rest and K^- interactions in flight. Thirty-seven Σ hyperons with decays have been observed. Of these, 26 were produced by K^- absorptions at rest, 11 were produced by in-flight interactions. The angular distribution of the space angle in the center of mass between the initial Σ -hyperon direction and the decay-pion direction was measured.

Figure 7 illustrates the resulting distribution, based on intervals of the cosine of the pion space angle of $1/3$, from -1 to $+1$. This corresponds to equal intervals of solid angle.

The distribution tends towards isotropy.

Σ^+ Branching Ratio

Of the 37 hyperons observed, 28 decayed at rest, thus giving a definite indication that they were Σ^+ (Σ^- interacts strongly when at rest). A Σ hyperon decaying in flight into a pion can be either Σ^+ or Σ^- . Of the 28 decays at rest, nine decayed via the $(p + \pi^0)$ mode, the other 19 via the $(n + \pi^+)$ mode. Detection of the $(n + \pi^+)$ mode in emulsion work is efficient only to about 90%. Also, only decays at rest can be reliably identified.




Fig. 7. Angular spectrum of decay pions from hyperons produced in K^- -absorption stars. (θ is space angle between projected direction of Σ and direction of decay pion (π^+ or π^0 , as case may be) in center-of-mass system).

The published results of both emulsion and bubble chamber groups are summarized thus:

<u>Group</u>	<u>$(p + \pi^0)$ mode</u>	<u>$(n + \pi^+)$ mode</u>
Goldhaber et al. [9]	9	19
White et al. [16]	4	8
Barkas et al. [17]	13	13
Fry et al. [20]	<u>26</u>	<u>20</u>
	52	60
Corrected for efficiency	52	~ 66
Alvarez et al.	<u>27</u>	<u>31</u>
	79	97
Over-all ratio: $\Sigma_p / \Sigma_{\pi^+} = 79/97 = 0.81 \pm 0.12$		

ACKNOWLEDGMENTS

This work was conducted under the tutelage of Professor Gerson Goldhaber. It is but a small part of the diverse and considerable efforts in emulsion research made possible through his insight and his direction of the very capable group he has established.

The comparable group of Dr. Sulamith Goldhaber and Dr. Warren Chupp shares equally in its contributions to these efforts.

Specific thanks go also to Mr. Joseph Lannutti and Dr. Louis Jauneau for much unselfishly given personal aid.

This work was done under the auspices of the U. S. Atomic Energy Commission.

REFERENCES

1. Table of Particles, revised October, 1956, UCRL
2. Edwin L. Iloff, Interactions and Lifetimes of K Mesons (thesis) UCRL-3605, Nov. 1956.
3. A. Pais, Phys. Rev. 86, 663 (1952).
4. M. Gell-Mann and A. Pais, Proceedings of the Glasgow Conference on Nuclear and Meson Physics, Pergamon Press, London, 1955; T. Nakano and K. Nishijima, Prog. Theor. Phys. 10, 581 (1953); M. Goldhaber, Phys. Rev. 101, 433 (1956); R.G. Sachs, Phys. Rev. 99, 1573 (1955).
5. From data supplied in a letter dated August 5, 1955, from C. Waller, Ilford Limited, to A. H. Rosenfeld.
6. As an example, Proposal for the Detection of Mass-500 Particle, W. Chupp and S. Goldhaber, Feb. 1957, (unpublished).
7. Goldhaber, Goldsack, and Lannutti, Method for Alignment of Stripped Nuclear Emulsions, UCRL-2928, March 1955.
8. Confirmative calculations made by L. Janneau, Berkeley, November 1956.
9. Webb, Iloff, Featherston, Chupp, Goldhaber, and Goldhaber, A Study of Negative K Mesons (to be published).
10. W. H. Barkas and D. M. Young, Emulsion Tables I. Heavy-Particle Functions, UCRL-2579 (Rev.), Sept. 1954.
11. Data from Iloff [2], which include three events from Livermore and six events from Harvard-MIT, as published in Phys. Rev. 102, 927 (1956).
12. Cork, Lambertson, Piccioni, and Wenzel, Lifetime and Hydrogen Cross Section of K^- Mesons, Phys. Rev. 106, 167 (1957).
13. Proceedings of the Sixth Rochester Conference on High-Energy Physics, 1956, Interscience, New York, 1956, Section VI.
14. Private communication from Dr. M. Ceccarelli of the Göttingen group, April 1957.
15. J. E. Lannutti, A Study of the Interaction of Positive K Mesons (thesis), University of California (to be published).
16. Gilbert, Violet, and White, Negative K-Particle Captures by Bound and Free Protons in Emulsion, UCRL-4814, Feb. 1957.

REFERENCES

1. Taylor, W. *Spencer's Grammar of the English Language*. 1900. (Oxford)
2. *English Grammar*. 1900. (Oxford)
3. *English Grammar*. 1900. (Oxford)
4. *English Grammar*. 1900. (Oxford)
5. *English Grammar*. 1900. (Oxford)
6. *English Grammar*. 1900. (Oxford)
7. *English Grammar*. 1900. (Oxford)
8. *English Grammar*. 1900. (Oxford)
9. *English Grammar*. 1900. (Oxford)
10. *English Grammar*. 1900. (Oxford)
11. *English Grammar*. 1900. (Oxford)
12. *English Grammar*. 1900. (Oxford)
13. *English Grammar*. 1900. (Oxford)
14. *English Grammar*. 1900. (Oxford)
15. *English Grammar*. 1900. (Oxford)
16. *English Grammar*. 1900. (Oxford)

17. Barkas, Dudziak, Giles, Heckman, Inman, Mason, Nichols, and Smith, Phys. Rev. 105, 1417 (1957)
18. T. D. Lee and C. N. Yang, Possible Interference Phenomena between Parity Doublets, Phys. Rev. (to be published).
19. Alvarez, Bradner, Falk-Vairant, Gow, Rosenfeld, Solmitz, and Tripp. K^- Interactions in Hydrogen, UCRL-3583, Nov. 1956.
20. Fry, Schneps, Snow, Swami, and Wold, Charged Σ Hyperons from 1001 K^- -Meson Stars, Department of Physics, University of Wisconsin, Madison, Wisconsin.

7 Sept. 59

INTERLIB
AEC

Thesis

35706

F2515

Featherston

Interactions of negative K mesons in flight in nuclear emulsion.

7 Sept. 59

INTERLIB
AEC

35706

Thesis


F2515

Featherston

Interactions of negative K mesons in flight in nuclear emulsion.

thesF2515

Interactions of negative K meson in flig



3 2768 002 13408 2

DUDLEY KNOX LIBRARY